

Subject: Music therapy for Psychiatry long-stay patients of Social Dysfunction intervention case report

Abstract: Mental patients in our country, Prolonged hospitalization and repeated hospitalization is relatively common. At the same time of clinical symptoms, there is still a social and cognitive impairment, affect the quality of life. Although antipsychotic drugs can treat hallucinations, delusions, but can't improve social skills. This article selects convalescence schizophrenia, depression and obsessive-compulsive disorder in 3 different conditions of prolonged hospitalization patients, Using Receptive Methods in Music Therapy and Recreative Music Therapy for social skills training theory and method of case report, to carry out the relevant treatment and research to provide the reference.

References:

- [1] (英) / T r o w e r P , / B r i M e d J 19 5 7 , 2 9 4 (1 4) - 6 6 3 - 6 6 4
- [2] 曹新妹. 实用精神科护理 [M]. 上海: 上海科学技术出版社. 2007. 3.
- [4] Bustillo JR;Lauriello J;Keith SJ Schizophrenia:improving outcome.1999(05)
- [5] Mueser K;Bellack A;Douglas M Prevalence and stability of social skill deficits in schizophrenia 1991(05)
- [6] 蒋锋, 汤宜郎, 侯也之. 精神疾病病耻感形成的相关因素及对 [J]. 中国心理卫生杂志, 2002, 16(10): 317- 319

Keyword: Music therapy, Psychiatric, Social skill training, Long-stay